

Spatiotemporal variability of snow properties derived from AVIRIS

Felix Seidel¹, Karl Rittger², Mckenzie Skiles³, Noah Molotch^{1,4}, Tom Painter¹



¹Jet Propulsion Laboratory / California Institute of Technology, Pasadena CA, USA

²National Snow and Ice Data Center, University of Colorado, Boulder CO, USA

³Earth Science Department, Utah Valley University, Orem, UT, USA

⁴Department of Geography, Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO, USA

This work was performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

©2017 All rights reserved



Jet Propulsion Laboratory
California Institute of Technology

The Cryosphere, 10, 1229–1244, 2016
www.the-cryosphere.net/10/1229/2016/
doi:10.5194/tc-10-1229-2016
© Author(s) 2016. CC Attribution 3.0 License.



Case study of spatial and temporal variability of snow cover, grain size, albedo and radiative forcing in the Sierra Nevada and Rocky Mountain snowpack derived from imaging spectroscopy

Felix C. Seidel¹, Karl Rittger², S. McKenzie Skiles¹, Noah P. Molotch^{3,1}, and Thomas H. Painter¹

¹Jet Propulsion Laboratory/California Institute of Technology, Pasadena, CA, USA

²National Snow and Ice Data Center, University of Colorado, Boulder, CO, USA

³Department of Geography, Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO, USA


Correspondence to: Felix C. Seidel (felix.c.seidel@gmail.com)

Sierra Nevada Study Area

Kings and Kaweah river basins

Sierra Nevada, CA. 2009


Dates	27 Feb., 05 Mar., 04 Apr., 07 May, 08 May
Pixel Size [m]	14.5
Image Area [km ² ,pixels]	252.03067, $1.2 \cdot 10^6$
Elevation Min./Mean/Max. [m]	1140 / 2575 / 3533



36.75

36.5

36.25




-118.75

JPL

-118.5

Snow Hydrology

Sierra Nevada study site



- Maritime snowpack (typically warmer and deeper than continental)
- 2009 was an average snow year in the Sierra Nevada

Retrieval Algorithms

Directional reflectance

$$HDRF_{sfc}^{obs} = \frac{R_{sensor}^{obs} - R_{atm}^{mdl}}{T_{\uparrow, mdl} + S_{sfc}^{mdl} (R_{sensor}^{obs} - R_{atm}^{mdl})}.$$



More information:

Painter, T. H., Seidel, F. C., Bryant, A. C., McKenzie Skiles, S., and Rittger, K.: Imaging spectroscopy of albedo and radiative forcing by light-absorbing impurities in mountain snow, *Journal of Geophysical Research: Atmospheres*, 118, 9511–9523, doi:10.1002/jgrd.50520, 2013.

Retrieval Algorithms

Directional reflectance

$$HDRF_{sfc}^{obs} = \frac{R_{sensor}^{obs} - R_{atm}^{mdl}}{T_{\uparrow, mdl} + S_{sfc}^{mdl} (R_{sensor}^{obs} - R_{atm}^{mdl})}.$$

Snow grain size

$$r : \left[\min \sum_{i=1.17 \mu m}^{1.27 \mu m} \left| HDRF_{snow}^{mdl}(\theta_{\odot}, \theta_v, \varphi; r; \lambda_i) - HDRF_{snow}^{obs}(\theta_{\odot}, \theta_v, \varphi; r; \lambda_i) \right| \right].$$



Retrieval Algorithms

Directional reflectance

$$HDRF_{sfc}^{obs} = \frac{R_{sensor}^{obs} - R_{atm}^{mdl}}{T_{\uparrow, mdl} + S_{sfc}^{mdl} (R_{sensor}^{obs} - R_{atm}^{mdl})}.$$

Snow grain size

$$r : \left[\min \sum_{i=1.17 \mu m}^{1.27 \mu m} \left| HDRF_{snow}^{mdl}(\theta_{\odot}, \theta_v, \varphi; r; \lambda_i) - HDRF_{snow}^{obs}(\theta_{\odot}, \theta_v, \varphi; r; \lambda_i) \right| \right].$$

Snow albedo

$$\alpha_{snow}^{obs}(r, \lambda) = \frac{HDRF_{snow}^{obs}(\theta_{\odot}, \theta_v, \varphi; r; \lambda) \alpha_{snow}^{mdl}(r, \lambda)}{HDRF_{snow}^{mdl}(\theta_{\odot}, \theta_v, \varphi; r; \lambda)}$$



Retrieval Algorithms

Directional reflectance

$$HDRF_{sfc}^{obs} = \frac{R_{sensor}^{obs} - R_{atm}^{mdl}}{T^{\uparrow, mdl} + S_{sfc}^{mdl} (R_{sensor}^{obs} - R_{atm}^{mdl})}.$$

Snow grain size

$$r : \left[\min \sum_{i=1.17 \mu m}^{1.27 \mu m} \left| HDRF_{snow}^{mdl}(\theta_{\odot}, \theta_v, \varphi; r; \lambda_i) - HDRF_{snow}^{obs}(\theta_{\odot}, \theta_v, \varphi; r; \lambda_i) \right| \right].$$


Snow albedo

$$\alpha_{snow}^{obs}(r, \lambda) = \frac{HDRF_{snow}^{obs}(\theta_{\odot}, \theta_v, \varphi; r; \lambda) \alpha_{snow}^{mdl}(r, \lambda)}{HDRF_{snow}^{mdl}(\theta_{\odot}, \theta_v, \varphi; r; \lambda)}$$


Snow radiative forcing by light absorbing impurities

$$RF_{snow} = \int_{\lambda=0.36 \mu m}^{1.08 \mu m} E_{sfc}^{mdl}(\lambda, \theta_{\odot}) \left[\alpha_{snow}^{mdl}(r, \lambda) - \alpha_{snow}^{obs}(r, \lambda) c(r, \lambda_c) \right] \Delta \lambda,$$

Time Series of Area Mean Values



Snow Grain Size




Albedo

27.Feb. 2009

04.Apr. 2009

08.May. 2009

Sierra Nevada, California, USA




Radiative Forcing

27.Feb. 2009


04.Apr. 2009

08.May. 2009

Sierra Nevada, California, USA




27.Feb. 2009




04.Apr. 2009

27.Feb. 2009




04.Apr. 2009


27.Feb. 2009




04.Apr. 2009




08.May. 2009




08.May. 2009



08.May. 2009




**Sierra Nevada
California, U.S.**



**Sierra Nevada
California, U.S.**

0.4






**Sierra Nevada
California, U.S.**




0 50

Grain Size,




Albedo, and Radiative Forcing v.s. Elevation






27 Feb 2009






05 Mar 2009



04 Apr 2009



07 May 2009



08 May 2009

50 100 150 200 250
Frequency [scaled 15 to 255]

50 100 150 200 255
Frequency [scaled 15 to 255]

50 100 150 200 250
Frequency [scaled 15 to 255]

Grain Size,

Albedo, and Radiative Forcing v.s. Terrain Aspect




27 Feb 2009

05 Mar 2009

04 Apr 2009

07 May 2009

08 May 2009



Conclusions

- Complex local/regional climate-snow-terrain feedbacks with snow
- ASO and this study are first to show spatially continuous snow grain size, albedo, and radiative forcing at multiple time steps during the ablation period thanks to imaging spectroscopy.
- Intuitive results:
 - smaller grain sizes on north facing slopes
 - inverse relationship between grain size and albedo
- New and counterintuitive results:
 - faster grain growth rates at higher elevations
 - higher grain sizes at high elevations during strong melt
- Our methods and algorithms can be applied to e.g. HypsIRI

Thank You!

Felix.Seidel@jpl.nasa.gov



Jet Propulsion Laboratory
California Institute of Technology

jpl.nasa.gov